



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PUBLICATIONS RECEIVED.

Birds of Springfield and Vicinity.—By Robert O. Morris. The town of Springfield lies on the Connecticut river but a few miles north of the dividing line between Massachusetts and Connecticut. In the preparation of the work the author passes over well beaten ground. From this very place the Rev. Wm. B. O. Peabody, Dr. J. A. Allen, F. H. Keys and W. W. Colburn, have enriched our ornithological literature; while Messrs. Emmons, Clark, Waters, Hoffman, and Drs. Merriam, Wood, and Faxon, have also done more or less work within the territory covered by the present volume. It has all appearance of having been done in a most thorough manner. The style is easy, direct and pleasing; many very valuable facts are related in an unassuming manner, and the description of the surrounding country is particularly fine and easily comprehended. 254 species and sub-species are recorded, all of which, with the exception of less than a fifth part, have come under the writer's personal observation. To the above he adds 4 species of probable occurrence but excluded from insufficient evidence; 5 introduced species, but one surviving—the ubiquitous and ever increasing European House Sparrow; and finally 2 extirpated species—the Heath Hen and Wild Turkey.

A work of this kind bespeaks long experience, and Mr. Morris is to be congratulated upon the fact that the elegance of his diction is further embellished by the work of the publisher.

[F. L. B.]

A Quantitative Study of Variation in the Smaller North-America Shrikes.—This quantitative study of variation to determine the status of a sub-species of birds, by our President, R. M. Strong is a pioneer in the application of an exact criterion for the determination of species and sub-species. There is no little objection to this quantitative method of study from one quarter and another, for no very good reason we can see. Manifestly no other kind of study can bring us exact results. A quantitative study is possible only when a large amount of material is available from many different regions, and this is